

Processus gaussiens pour l'exploration d'espaces de paramètres. De l'optimisation à l'estimation conservative d'ensembles d'excursion

David Ginsbourger¹

Acknowledgements: a number of co-authors, notably appearing via citations!

¹IMSV, Department of Mathematics and Statistics, University of Bern

Inauguration du GIS LARTISSTE

18 Octobre 2022

Preamble

While statistics is increasingly employed to address societal needs, there is a number in situations where data (un)availability is a potential bottleneck:

- Environmental hazards, especially in a climate change context
- Safety of industrial installations such as power plants
- Personalized/precision medicine
- Exploration of natural resources, surveys, etc.

Preamble

While statistics is increasingly employed to address societal needs, there is a number in situations where data (un)availability is a potential bottleneck:

- Environmental hazards, especially in a climate change context
- Safety of industrial installations such as power plants
- Personalized/precision medicine
- Exploration of natural resources, surveys, etc.

It is not always that the absolute amount of data is moderate, but rather that useful data for the question at hand is rare, expensive, confidential. . .

Preamble

Numerical simulations (also known as “Computer Experiments”) are more and more used to cope with the lack of observational data. Yet they also come with issues as accurate simulations can be quite resource-intensive!

Preamble

[Numerical simulations](#) (also known as “Computer Experiments”) are more and more used to cope with the lack of observational data. Yet they also come with issues as accurate simulations can be quite resource-intensive!

Models inherited from [spatial statistics](#) turn out to be of great use in [uncertainty quantification](#) to deal with both observational and simulation data.

Preamble

[Numerical simulations](#) (also known as “Computer Experiments”) are more and more used to cope with the lack of observational data. Yet they also come with issues as accurate simulations can be quite resource-intensive!

Models inherited from [spatial statistics](#) turn out to be of great use in [uncertainty quantification](#) to deal with both observational and simulation data.

In particular, the former offer a neat framework for the definition of [sequential design](#) strategies for “uncertainty reduction”, or, in other words, data acquisition policies dedicated to address specific research questions.

Outline

- 1 Introduction
 - Settings
 - From GP-based optimization to set estimation
- 2 Focus on (reducing) uncertainties on sets
 - Introduction to Sequential Uncertainty Reduction strategies
 - Towards conservative excursion set estimation

Set up of main contributions presented next

We investigate i) a complex system represented by a deterministic function $f : \mathbf{x} \in E \mapsto f(\mathbf{x}) \in F$, ii) and/or quantities relying on f , based on a limited number of evaluations of f .

Set up of main contributions presented next

We investigate i) a complex system represented by a deterministic function $f : \mathbf{x} \in E \mapsto f(\mathbf{x}) \in F$, ii) and/or quantities relying on f , based on a limited number of evaluations of f .

Two typical examples

- **Safety engineering**: \mathbf{x} is a vector parametrizing some system and f returns an indicator of dangerousness. It is then crucial to understand which \mathbf{x} 's lead to "high" values of $f(\mathbf{x})$.

Set up of main contributions presented next

We investigate i) a complex system represented by a deterministic function $f : \mathbf{x} \in E \mapsto f(\mathbf{x}) \in F$, ii) and/or quantities relying on f , based on a limited number of evaluations of f .

Two typical examples

- **Safety engineering**: \mathbf{x} is a vector parametrizing some system and f returns an indicator of dangerousness. It is then crucial to understand which \mathbf{x} 's lead to “high” values of $f(\mathbf{x})$.
- **Flow dynamics**: \mathbf{x} stands e.g. for the medium, boundary conditions, etc. and f returns the evolution of a fluid and/or a measure of discrepancy between simulation results and given observation results.

Function approximation by means of GP models

Typical situation : f was evaluated at a set of “points” $\mathbf{x}_1, \dots, \mathbf{x}_n \in D \subset E$ and one wishes to estimate a quantity relying on f and/or perform new evaluations in order to improve this estimation.

Function approximation by means of GP models

Typical situation : f was evaluated at a set of “points” $\mathbf{x}_1, \dots, \mathbf{x}_n \in D \subset E$ and one wishes to estimate a quantity relying on f and/or perform new evaluations in order to improve this estimation.

\Rightarrow legitimate to rely on some approximation(s) of f knowing $f(\mathbf{x}_i) + \epsilon_i$ ($1 \leq i \leq n$). A number of approaches do exist. . .

Function approximation by means of GP models

Typical situation : f was evaluated at a set of “points” $\mathbf{x}_1, \dots, \mathbf{x}_n \in D \subset E$ and one wishes to estimate a quantity relying on f and/or perform new evaluations in order to improve this estimation.

\Rightarrow legitimate to rely on some approximation(s) of f knowing $f(\mathbf{x}_i) + \epsilon_i$ ($1 \leq i \leq n$). A number of approaches do exist. . .

Principles of the [Gaussian Process approach](#) (GP): suppose that, *a priori*, f is a realization of a GP $(Z(\mathbf{x}))_{\mathbf{x} \in D}$ and approximate f and/or the quantities of interest via the **conditional distribution** of Z knowing $Z(\mathbf{x}_i) + \epsilon_i = f(\mathbf{x}_i) + \epsilon_i$.

Function approximation by means of GP models

Typical situation : f was evaluated at a set of “points” $\mathbf{x}_1, \dots, \mathbf{x}_n \in D \subset E$ and one wishes to estimate a quantity relying on f and/or perform new evaluations in order to improve this estimation.

\Rightarrow legitimate to rely on some approximation(s) of f knowing $f(\mathbf{x}_i) + \epsilon_i$ ($1 \leq i \leq n$). A number of approaches do exist. . .


Principles of the **Gaussian Process approach** (GP): suppose that, *a priori*, f is a realization of a GP $(Z(\mathbf{x}))_{\mathbf{x} \in D}$ and approximate f and/or the quantities of interest via the **conditional distribution** of Z knowing $Z(\mathbf{x}_i) + \epsilon_i = f(\mathbf{x}_i) + \epsilon_i$.

\Rightarrow very practical for **sequential design of experiments**.

Example inverse problem in hydrogeology

A costly full factorial experimental design!

Misfit (objective function)



Application of GP-based optimization

The previous example was produced in the framework of a collaboration with [G. Pirot](#) (University of Lausanne), [T. Kityakierne](#) (now at Mahidol University, Bangkok) and [P. Renard](#) (University of Neuchâtel).

⇒ See recently published Hydrol. Earth Syst. Sci. paper (2019).

The previous example was produced in the framework of a collaboration with [G. Pirot](#) (University of Lausanne), [T. Kityakierne](#) (now at Mahidol University, Bangkok) and [P. Renard](#) (University of Neuchâtel).

⇒ See recently published Hydrol. Earth Syst. Sci. paper (2019).

Main focus in most of the following

In a related set-up, how to estimate parameter [regions](#) where f takes target values using such models and dedicated sequential design strategies?

The previous example was produced in the framework of a collaboration with [G. Pirot](#) (University of Lausanne), [T. Kityakierne](#) (now at Mahidol University, Bangkok) and [P. Renard](#) (University of Neuchâtel).

⇒ See recently published Hydrol. Earth Syst. Sci. paper (2019).

Main focus in most of the following

In a related set-up, how to estimate parameter [regions](#) where f takes target values using such models and dedicated sequential design strategies?

As a transition, let us review a few selected seminal references about GP modelling and GP-based “Bayesian Optimization”.

A few references on GP modelling . . .



A. O'Hagan (1978).

Curve fitting and optimal design for prediction.

Journal of the Royal Statistical Society, Series B, 40(1):1-42.



J. Sacks, W.J. Welch, T.J. Mitchell, and H. P. Wynn (1989).

Design and Analysis of Computer Experiments

Statist. Sci. 4(4), 409-423.



H. Omre and K. Halvorsen (1989).

The bayesian bridge between simple and universal kriging.

Mathematical Geology, 22 (7):767-786.



M. S. Handcock and M. L. Stein (1993).

A bayesian analysis of kriging.

Technometrics, 35(4):403-410.



A.W. Van der Vaart and J. H. Van Zanten (2008).

Rates of contraction of posterior distributions based on Gaussian process priors.

Annals of Statistics, 36:1435-1463.

... and on GP-based Optimization



H.J. Kushner (1964).

A new method of locating the maximum of an arbitrary multi-peak curve in the presence of noise.
Journal of Basic Engineering, 86:97-106.



J. Mockus (1972).

On Bayesian methods for seeking the extremum.
Automatics and Computers (Avtomatika i Vychislitel'naya Tekhnika), 4(1):53-62.



J. Mockus, V. Tiesis, and A. Zilinskas (1978).

The application of Bayesian methods for seeking the extremum.
In Dixon, L. C. W. and Szegő, G. P., editors, *Towards Global Optimisation*, volume 2, pages 117-129. Elsevier Science Ltd., North Holland, Amsterdam.



J.M. Calvin (1997).

Average performance of a class of adaptive algorithms for global optimization.
The Annals of Applied Probability, 7(3):711-730.



M. Schonlau, W.J. Welch and D.R. Jones (1998).

Efficient Global Optimization of Expensive Black-box Functions.
Journal of Global Optimization.

Outline

1 Introduction

- Settings
- From GP-based optimization to set estimation

2 Focus on (reducing) uncertainties on sets

- Introduction to Sequential Uncertainty Reduction strategies
- Towards conservative excursion set estimation

Our main topic today: background and motivations

A number of practical problems boil down to determining sets of the form

$$I^* = \{\mathbf{x} \in D : f(\mathbf{x}) \in T\} = f^{-1}(T)$$

where $f : D \rightarrow \mathbb{R}^k$ ($k \geq 1$), $D \subset \mathbb{R}^d$ ($d \geq 1$), and $T \subset \mathbb{R}^k$.

Our main topic today: background and motivations

A number of practical problems boil down to determining sets of the form

$$\Gamma^* = \{\mathbf{x} \in D : f(\mathbf{x}) \in T\} = f^{-1}(T)$$

where $f : D \rightarrow \mathbb{R}^k$ ($k \geq 1$), $D \subset \mathbb{R}^d$ ($d \geq 1$), and $T \subset \mathbb{R}^k$.

Examples

- Contour lines
- Excursion/sojourn sets above/below thresholds
- Admissible regions in constrained optimization
- High gradient/high curvature regions, etc.
- (Pareto sets in multi-objective optimization... but then T depends on f !)

Background and motivations

We essentially focus today on the case where $k = 1$, D is compact, f is continuous, and $T = [t, +\infty)$ or $(-\infty, t]$ for some prescribed $t \in \mathbb{R}$.

$\Gamma^* = \{\mathbf{x} \in D : f(\mathbf{x}) \geq t\}$ is then referred to as the **excursion set of f above t** .

Background and motivations

We essentially focus today on the case where $k = 1$, D is compact, f is continuous, and $T = [t, +\infty)$ or $(-\infty, t]$ for some prescribed $t \in \mathbb{R}$.

$\Gamma^* = \{\mathbf{x} \in D : f(\mathbf{x}) \geq t\}$ is then referred to as the **excursion set of f above t** .

Our aim is to estimate Γ^* and quantify uncertainty on it when f can solely be evaluated at a few points, both in static and sequential cases.

Test case from safety engineering




Figure: Excursion set (light gray) of a nuclear criticality safety coefficient depending on two design parameters. Blue triangles: initial experiments.



C. Chevalier (2013).

Fast uncertainty reduction strategies relying on Gaussian process models.

Ph.D. thesis, University of Bern.

Making a sensible estimation of I^* based on a drastically limited number of evaluations $f(\mathbf{X}_n) = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_n))'$ calls for additional assumptions on f .

Making a sensible estimation of I^* based on a drastically limited number of evaluations $f(\mathbf{X}_n) = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_n))'$ calls for additional assumptions on f .

As before, we consider the Bayesian framework where a Gaussian Process (GP) prior is put on f , i.e. f is seen as one realization of a GP $(Z(\mathbf{x}))_{\mathbf{x} \in D}$ (characterized in distribution by a mean m and a covariance kernel k).

Making a sensible estimation of Γ^* based on a drastically limited number of evaluations $f(\mathbf{X}_n) = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_n))'$ calls for additional assumptions on f .

As before, we consider the Bayesian framework where a Gaussian Process (GP) prior is put on f , i.e. f is seen as one realization of a GP $(Z(\mathbf{x}))_{\mathbf{x} \in D}$ (characterized in distribution by a mean m and a covariance kernel k).

In the GP set-up, the main object of interest is represented by

$$\Gamma = \{\mathbf{x} \in D : Z(\mathbf{x}) \in T\} = Z^{-1}(T)$$

Under our previous assumptions on T and assuming that is chosen Z with continuous paths, Γ is a **Random Closed Set** (See thesis below for detail).



D. Azzimonti (2016).

Contributions to Bayesian set estimation relying on random field priors.

Ph.D. thesis, University of Bern.

Simulating excursion sets under a GRF model

Posterior simulations on a 50×50 grid of Z and Γ knowing $Z(\mathbf{X}_n) = f(\mathbf{X}_n)$.

How to quantify the uncertainty on Γ ?

There are many ways to quantify uncertainties on sets!

How to quantify the uncertainty on Γ ?

There are many ways to quantify uncertainties on sets!

This will be one of the recurring questions throughout the talk, but we will not be exhaustive by far. For more detail see, e.g.,



I. Molchanov (2005)

Theory of Random Sets.

Springer.



D. Azzimonti, J. Bect, C. Chevalier and D. Ginsbourger (2016).

Quantifying uncertainties on excursion sets under a Gaussian random field prior.

SIAM/ASA Journal on Uncertainty Quantification.

How to quantify the uncertainty on Γ ?

There are many ways to quantify uncertainties on sets!

This will be one of the recurring questions throughout the talk, but we will not be exhaustive by far. For more detail see, e.g.,



I. Molchanov (2005)

Theory of Random Sets.

Springer.



D. Azzimonti, J. Bect, C. Chevalier and D. Ginsbourger (2016).


Quantifying uncertainties on excursion sets under a Gaussian random field prior.

SIAM/ASA Journal on Uncertainty Quantification.


Before moving to random set-related concepts, a first spontaneous idea is to “scalarize” the problem, for instance by looking at Γ 's volume. Let us make a detour through some GP basics in order to do so.

Kriging (Gaussian Process Interpolation)

Posterior mean




Posterior variance



$$\begin{cases} m_n(\mathbf{x}) = m(\mathbf{x}) + k(\mathbf{X}_n, \mathbf{x})^T k(\mathbf{X}_n, \mathbf{X}_n)^{-1} (f(\mathbf{X}_n) - m(\mathbf{X}_n)) \\ s_n^2(\mathbf{x}) = k(\mathbf{x}, \mathbf{x}) - k(\mathbf{X}_n, \mathbf{x})^T k(\mathbf{X}_n, \mathbf{X}_n)^{-1} k(\mathbf{X}_n, \mathbf{x}) \end{cases}$$

Conditional probability of excursion



From $\mathcal{L}_n(Z(\mathbf{x})) = \mathcal{N}(m_n(\mathbf{x}), s_n^2(\mathbf{x}))$, the “coverage probability” of Γ (or conditional/posterior probability of excursion, here) can be expanded as

$$p_n(\mathbf{x}) = \mathbb{P}_n(\mathbf{x} \in \Gamma) = \mathbb{P}_n(Z(\mathbf{x}) \geq t) = \Phi\left(\frac{m_n(\mathbf{x}) - t}{s_n(\mathbf{x})}\right)$$

Outline

- 1 Introduction
 - Settings
 - From GP-based optimization to set estimation
- 2 Focus on (reducing) uncertainties on sets
 - Introduction to Sequential Uncertainty Reduction strategies
 - Towards conservative excursion set estimation

From p_n to moments of Γ 's volume

Denote by μ a finite measure on $(D, \mathcal{B}(D))$ [one can think here of μ as Vol] and set $\alpha^* = \mu(\Gamma^*)$, i.e. the “volume of excursion” in the considered case.

The GP model leads to a random analogue $\alpha = \mu(\Gamma)$, and by Robbins' theorem, the posterior expectation of α can be written in terms of p_n :

$$\mathbb{E}_n[\mu(\Gamma)] = \mathbb{E}_n \left[\int_D \mathbf{1}_\Gamma(\mathbf{u}) d\mu(\mathbf{u}) \right] = \int_D p_n(\mathbf{u}) d\mu(\mathbf{u})$$

From p_n to moments of Γ 's volume

Denote by μ a finite measure on $(D, \mathcal{B}(D))$ [one can think here of μ as Vol] and set $\alpha^* = \mu(\Gamma^*)$, i.e. the “volume of excursion” in the considered case.

The GP model leads to a random analogue $\alpha = \mu(\Gamma)$, and by Robbins' theorem, the posterior expectation of α can be written in terms of p_n :

$$\mathbb{E}_n[\mu(\Gamma)] = \mathbb{E}_n \left[\int_D \mathbf{1}_\Gamma(\mathbf{u}) d\mu(\mathbf{u}) \right] = \int_D p_n(\mathbf{u}) d\mu(\mathbf{u})$$

However, the (posterior) distribution of α has been considered analytically intractable.



R.J. Adler (2000)

On excursion sets, tube formulas and maxima of random fields.
Annals of Applied Probability, 10(1):1-74.



E. Vazquez and M. Piera Martinez (2006).

Estimation of the volume of an excursion set of a Gaussian process using intrinsic Kriging.

arXiv:math/0611273 [math.ST].

About conditional moments of α

Fortunately, as already pointed out in Molchanov 2005 in more general settings, $\mathbb{E}_n[\alpha^r]$ can also be worked out for $r \geq 2$, at the price of calculating integrals. In our framework, we have indeed:

$$\begin{aligned}
 \mathbb{E}_n[\alpha^r] &= \mathbb{E}_n \left[\left(\int_D \mathbf{1}_\Gamma(\mathbf{u}) d\mu(\mathbf{u}) \right)^r \right] \\
 &= \mathbb{E}_n \left[\left(\int_D \mathbf{1}_\Gamma(\mathbf{u}_1) d\mu(\mathbf{u}_1) \right) \dots \left(\int_D \mathbf{1}_\Gamma(\mathbf{u}_r) d\mu(\mathbf{u}_r) \right) \right] \\
 &= \int_D \dots \int_D \mathbb{E}_n[\mathbf{1}_\Gamma(\mathbf{u}_1) \dots \mathbf{1}_\Gamma(\mathbf{u}_r)] d\mu(\mathbf{u}_1) \dots d\mu(\mathbf{u}_r) \\
 &= \int_D \dots \int_D \mathbb{P}_n(Z(\mathbf{u}_1) \geq t, \dots, Z(\mathbf{u}_r) \geq t) d\mu(\mathbf{u}_1) \dots d\mu(\mathbf{u}_r)
 \end{aligned}$$

Hence, recalling the GP assumption, $\mathbb{E}_n[\alpha^r]$ writes as an r -dimensional integral which integrand involves a **r -dimensional Gaussian CDF**.

A useful bound for the case $r = 2$

In what follows, the case $r = 2$ will be of special importance as we will consider sequential design strategies aiming at reducing $\text{Var}_n[\alpha]$.

The following underlined quantity, that is easier to compute and also comes with a nice interpretation, has been used as well:

$$\begin{aligned}
 \text{Var}_n[\alpha] &= \mathbb{E}_n \left[\left(\int_D (\mathbf{1}_\Gamma(\mathbf{u}) - p_n(\mathbf{u})) d\mu(\mathbf{u}) \right)^2 \right] \\
 &\leq \mu(D)^2 \mathbb{E}_n \left[\int_D (\mathbf{1}_\Gamma(\mathbf{u}) - p_n(\mathbf{u}))^2 d\mu(\mathbf{u}) \right] \\
 &= \mu(D)^2 \underbrace{\int_D p_n(\mathbf{u})(1 - p_n(\mathbf{u})) d\mu(\mathbf{u})}_{\text{Integrated indicator variance}}
 \end{aligned}$$

A useful bound for the case $r = 2$

In what follows, the case $r = 2$ will be of special importance as we will consider sequential design strategies aiming at reducing $\text{Var}_n[\alpha]$.

The following underlined quantity, that is easier to compute and also comes with a nice interpretation, has been used as well:

$$\begin{aligned} \text{Var}_n[\alpha] &= \mathbb{E}_n \left[\left(\int_D (\mathbf{1}_\Gamma(\mathbf{u}) - p_n(\mathbf{u})) d\mu(\mathbf{u}) \right)^2 \right] \\ &\leq \mu(D)^2 \mathbb{E}_n \left[\int_D (\mathbf{1}_\Gamma(\mathbf{u}) - p_n(\mathbf{u}))^2 d\mu(\mathbf{u}) \right] \\ &= \mu(D)^2 \underbrace{\int_D p_n(\mathbf{u})(1 - p_n(\mathbf{u})) d\mu(\mathbf{u})}_{\text{Integrated indicator variance}} \end{aligned}$$

The excursion volume's variance and the integrated indicator variance are used as two particular “measures of uncertainty” in what follows.

Towards Stepwise Uncertainty Reduction strategies

Let us informally consider the following **1-step-lookahead** scheme:

- For some chosen (say, non-negative) functional defined on GP distributions, define the **uncertainty at time $n \geq 0$** , H_n , as this functional applied to the current posterior GP (E.g., $H_n = \text{var}_n(\alpha)$).
- Starting from some initial design $\{\mathbf{x}_1, \dots, \mathbf{x}_{n_0}\}$, at each iteration $n \geq n_0$, evaluate f at a point \mathbf{x}_{n+1}^* minimizing the so-called **SUR criterion** associated with the chosen notion of uncertainty:

$$J_n(\mathbf{x}_{n+1}) := \mathbb{E}_n(H_{n+1}(\mathbf{x}_{n+1}))$$

Towards Stepwise Uncertainty Reduction strategies

Let us informally consider the following **1-step-lookahead** scheme:

- For some chosen (say, non-negative) functional defined on GP distributions, define the **uncertainty at time $n \geq 0$** , H_n , as this functional applied to the current posterior GP (E.g., $H_n = \text{var}_n(\alpha)$).
- Starting from some initial design $\{\mathbf{x}_1, \dots, \mathbf{x}_{n_0}\}$, at each iteration $n \geq n_0$, evaluate f at a point \mathbf{x}_{n+1}^* minimizing the so-called **SUR criterion** associated with the chosen notion of uncertainty:

$$J_n(\mathbf{x}_{n+1}) := \mathbb{E}_n(H_{n+1}(\mathbf{x}_{n+1}))$$

See notably the following paper and seminal references therein:



J. Bect, D. Ginsbourger, L. Li, V. Picheny and E. Vazquez.

Sequential design of computer experiments for the estimation of a probability of failure.

Statistics and Computing, 22(3):773-793, 2012.

SUR strategies: Two candidate uncertainties

Two possible definitions for the uncertainty H_n are considered below:

$$H_n := \text{Var}_n(\alpha)$$

$$\tilde{H}_n := \int_D p_n(1 - p_n) d\mu$$

SUR strategies: Two candidate uncertainties

Two possible definitions for the uncertainty H_n are considered below:

$$H_n := \text{Var}_n(\alpha)$$

$$\tilde{H}_n := \int_D p_n(1 - p_n) d\mu$$

Uncertainties:

$$H_n := \text{Var}_n(\alpha)$$

$$\tilde{H}_n := \int_{\mathbb{X}} p_n(1 - p_n) d\mu$$

SUR criteria:

$$J_n(\mathbf{x}) := \mathbb{E}_n(\text{Var}_{n+1}(\alpha))$$

$$\tilde{J}_n(\mathbf{x}) := \mathbb{E}_n \left(\int_D p_{n+1}(1 - p_{n+1}) d\mu \right)$$

Main challenge to calculate $\tilde{J}_n(\mathbf{x})$ (similar for $J_n(\mathbf{x})$): Obtain a closed form expression for $\mathbb{E}_n(p_{n+1}(1 - p_{n+1}))$ and integrate it.

Deriving SUR criteria

Proposition

$$\mathbb{E}_n(p_{n+1}(\mathbf{x})(1 - p_{n+1}(\mathbf{x}))) = \Phi_2 \left(\begin{pmatrix} a(\mathbf{x}) \\ -a(\mathbf{x}) \end{pmatrix}, \begin{pmatrix} c(\mathbf{x}) & 1 - c(\mathbf{x}) \\ 1 - c(\mathbf{x}) & c(\mathbf{x}) \end{pmatrix} \right)$$

- $\Phi_2(\cdot, M)$: c.d.f. of centred bivariate Gaussian with covariance matrix M
- $a(\mathbf{x}) := (m_n(\mathbf{x}) - t)/s_{n+1}(\mathbf{x})$,
- $c(\mathbf{x}) := s_n^2(\mathbf{x})/s_{n+1}^2(\mathbf{x})$



C. Chevalier, J. Bect, D. Ginsbourger, V. Picheny, E. Vazquez and Y. Richet.
Fast parallel kriging-based stepwise uncertainty reduction with application to the identification of an excursion set.

Technometrics, 56(4):455-465, 2014.

Deriving SUR criteria

Proposition

$$\mathbb{E}_n(p_{n+1}(\mathbf{x})(1 - p_{n+1}(\mathbf{x}))) = \Phi_2 \left(\begin{pmatrix} a(\mathbf{x}) \\ -a(\mathbf{x}) \end{pmatrix}, \begin{pmatrix} c(\mathbf{x}) & 1 - c(\mathbf{x}) \\ 1 - c(\mathbf{x}) & c(\mathbf{x}) \end{pmatrix} \right)$$

- $\Phi_2(\cdot, M)$: c.d.f. of centred bivariate Gaussian with covariance matrix M
- $a(\mathbf{x}) := (m_n(\mathbf{x}) - t)/s_{n+1}(\mathbf{x})$,
- $c(\mathbf{x}) := s_n^2(\mathbf{x})/s_{n+1}^2(\mathbf{x})$



C. Chevalier, J. Bect, D. Ginsbourger, V. Picheny, E. Vazquez and Y. Richet.
Fast parallel kriging-based stepwise uncertainty reduction with application to the identification of an excursion set.

Technometrics, 56(4):455-465, 2014.



C. Chevalier, V. Picheny and D. Ginsbourger.

The KrigInv package: An efficient and user-friendly R implementation of Kriging-based inversion algorithms.

Computational Statistics & Data Analysis, 71:1021-1034, 2014



Back to the test case with SUR

Batch-sequential SUR strategies




Figure: 3 SUR iterations (\tilde{J}_n criterion with $q = 4$)

About the consistency



J. Bect, F. Bachoc and D. Ginsbourger (2019).

A supermartingale approach to Gaussian process based sequential design of experiments.

Bernoulli 25 (4A), 2883-2919.

Further questions about SUR and UQ on sets

Of course, in operational conditions, asymptotic results are worthwhile. However, concrete finite-sample outputs such as estimates of Γ^* and quantifications of the associated uncertainty are required as well.

Further questions about SUR and UQ on sets

Of course, in operational conditions, asymptotic results are worthwhile. However, concrete finite-sample outputs such as estimates of Γ^* and quantifications of the associated uncertainty are required as well.

Now, n being fixed, how to estimate Γ^* and to assess/represent the variability of the corresponding estimate(s)?

Outline

- 1 Introduction
 - Settings
 - From GP-based optimization to set estimation
- 2 Focus on (reducing) uncertainties on sets
 - Introduction to Sequential Uncertainty Reduction strategies
 - Towards conservative excursion set estimation


How to summarize the posterior distribution of sets?

For application purposes, let us reverse the perspective and focus on the sojourn/excursion case **below** t , where $\Gamma = \{\mathbf{x} \in D : Z(\mathbf{x}) \leq t\}$ [say a “safe set”] and $p_n : \mathbf{x} \in D \rightarrow p_n(\mathbf{x}) = P_n(Z(\mathbf{x}) \leq t)$.

How to summarize the posterior distribution of sets?

For application purposes, let us reverse the perspective and focus on the sojourn/excursion case **below** t , where $\Gamma = \{\mathbf{x} \in D : Z(\mathbf{x}) \leq t\}$ [say a “safe set”] and $p_n : \mathbf{x} \in D \rightarrow p_n(\mathbf{x}) = P_n(Z(\mathbf{x}) \leq t)$.

Coverage probability function



Define the (conditional) quantiles of Γ as ρ -level sets of p_n :


$$\begin{aligned} Q_\rho &:= \{\mathbf{x} \in D : p_n(\mathbf{x}) \geq \rho\} \\ &= \{\mathbf{x} \in D : P_n(Z(\mathbf{x}) \leq t) \geq \rho\}. \end{aligned}$$

How well Q_ρ estimates Γ can be quantified for instance through the “expected deviation”:

$$\mathbb{E}_n(\mu(Q_\rho \Delta \Gamma))$$

Estimates of Γ^* : the Vorob'ev expectation

Coverage probability function (Vorob'ev expectation)



The **Vorob'ev expectation** of $\Gamma \mid (Z(\mathbf{x}_1) = f(x_1), \dots, Z(\mathbf{x}_n) = f(x_n))$ is the ρ^* level set of ρ_n such that

$$\mu(Q_{\rho^*}) = \mathbb{E}_n[\mu(\Gamma)].$$

It is a state of the art result that Q_{ρ^*} minimizes $S \rightarrow \mathbb{E}_n(\mu(S \Delta \Gamma))$ among all closed sets $S \subset \mathbb{R}^d$ with volume $\mathbb{E}_n[\mu(\Gamma)]$.



C. Chevalier, D. Ginsbourger, J. Bect, and Molchanov, I.

Estimating and quantifying uncertainties on level sets using the Vorob'ev expectation and deviation with Gaussian process models.


mODa 10 Advances in Model-Oriented Design and Analysis, Physica-Verlag HD, 2013



Estimates of Γ^* : some limitations of Q_ρ quantiles

In practice one often wish to give **confidence statements** on the estimates.

Coverage probability function (Level set 0.95)




Q_ρ contains points which have marginal probability at least ρ of being in Γ .

\Rightarrow no confidence statement on the probability of the actual excursion set containing this specific estimate.

Estimates of Γ^* : some limitations of Q_ρ quantiles

In practice one often wish to give **confidence statements** on the estimates.

Coverage probability function (Level set 0.95)



Q_ρ contains points which have marginal probability at least ρ of being in Γ .

\Rightarrow no confidence statement on the probability of the actual excursion set containing this specific estimate.

E.g., the probabilities of Q_ρ containing the excursion set (computed on a grid) are

- 0.67 for $\rho = 0.95$
- 0.009 for $\rho = 0.5$
- 0.019 for $\rho = 0.56$ (Vorob'ev)

Conservative Estimates of Γ^*

We denote by **conservative estimate** for

$\Gamma \mid (Z(\mathbf{x}_1) = f(x_1), \dots, Z(\mathbf{x}_n) = f(x_n))$ at level β the largest Q_ρ such that $\mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta$:

$$E_{t,\alpha} = \arg \max_{Q_\rho} \{\mu(Q_\rho) : \mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta\}$$



D. Bolin, F. Lindgren.

Excursion and contour uncertainty regions for latent Gaussian models.

Journal of the Royal Statistical Society: Series B (Statistical Methodology), 2014.

Conservative Estimates of Γ^*

We denote by **conservative estimate** for

$\Gamma \mid (Z(\mathbf{x}_1) = f(x_1), \dots, Z(\mathbf{x}_n) = f(x_n))$ at level β the largest Q_ρ such that $\mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta$:

$$E_{t,\alpha} = \arg \max_{Q_\rho} \{\mu(Q_\rho) : \mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta\}$$



D. Bolin, F. Lindgren.

Excursion and contour uncertainty regions for latent Gaussian models.

Journal of the Royal Statistical Society: Series B (Statistical Methodology), 2014.

Such conservative estimate $E_{t,\beta}$ is hence

- the largest quantile such that, with probability β , the response is below the threshold **simultaneously at each of its locations**.
- based on a confidence statement on the whole set

Computing conservative estimates

The computation of a conservative estimate

$$E_{t,\beta} = \arg \max_{Q_\rho} \{\mu(Q_\rho) : \mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta\}$$

presents two (nested) computational bottlenecks:

- 1 find the set with the maximum volume;
- 2 compute $\mathbb{P}_n(Q_\rho \subset \Gamma)$.

Computing conservative estimates

The computation of a conservative estimate

$$E_{t,\beta} = \arg \max_{Q_\rho} \{\mu(Q_\rho) : \mathbb{P}_n(Q_\rho \subset \Gamma) \geq \beta\}$$

presents two (nested) computational bottlenecks:

- 1 find the set with the maximum volume;
- 2 compute $\mathbb{P}_n(Q_\rho \subset \Gamma)$.

For recent work on computing the last term, see for instance



D. Azzimonti and D. Ginsbourger (2018).

Estimating orthant probabilities of high dimensional Gaussian vectors with an application to set estimation.

Journal of Computational and Graphical Statistics, 27:2, 255-267

Computing $\mathbb{P}_n(Q_\rho \subset \Gamma)$

If Q_ρ is discretized over a grid $W = \{w_1, \dots, w_m\}$, then

$$\mathbb{P}_n(Q_\rho \subset \Gamma) = \mathbb{P}_n(Z(\mathbf{w}_1) \leq t, \dots, Z(\mathbf{w}_m) \leq t) = 1 - \mathbb{P}_n\left(\max_{i=1, \dots, m} Z(\mathbf{w}_i) > t\right)$$

Computing $\mathbb{P}_n(Q_\rho \subset \Gamma)$

If Q_ρ is discretized over a grid $W = \{\mathbf{w}_1, \dots, \mathbf{w}_m\}$, then

$$\mathbb{P}_n(Q_\rho \subset \Gamma) = \mathbb{P}_n(Z(\mathbf{w}_1) \leq t, \dots, Z(\mathbf{w}_m) \leq t) = 1 - \mathbb{P}_n\left(\max_{i=1, \dots, m} Z(\mathbf{w}_i) > t\right)$$

There exists a number of algorithms to estimate

$\mathbb{P}_n(Z(\mathbf{w}_1) \leq t, \dots, Z(\mathbf{w}_m) \leq t)$:

- 1 quasi-MC integration techniques
 - very fast and reliable in small dimensions;
 - hardly usable for dimensions higher than 1000.
- 2 pure MC techniques:
 - dimension independent;
 - high number of simulations for small variance.

Computing $\mathbb{P}_n(Q_\rho \subset \Gamma)$

If Q_ρ is discretized over a grid $W = \{\mathbf{w}_1, \dots, \mathbf{w}_m\}$, then

$$\mathbb{P}_n(Q_\rho \subset \Gamma) = \mathbb{P}_n(Z(\mathbf{w}_1) \leq t, \dots, Z(\mathbf{w}_m) \leq t) = 1 - \mathbb{P}_n\left(\max_{i=1, \dots, m} Z(\mathbf{w}_i) > t\right)$$

There exists a number of algorithms to estimate

$\mathbb{P}_n(Z(\mathbf{w}_1) \leq t, \dots, Z(\mathbf{w}_m) \leq t)$:

- 1 quasi-MC integration techniques
 - very fast and reliable in small dimensions;
 - hardly usable for dimensions higher than 1000.
- 2 pure MC techniques:
 - dimension independent;
 - high number of simulations for small variance.

IRSN test case

- an estimate with a good resolution requires an 100×100 grid for D ;
- W consists of +1000 grid points for some Q_ρ .

$\mathbb{P}_n(\max_{W \in W} Z_W > T)$: proposed hybrid algorithm

Algorithm:

- ① select q grid points, denoted $W_q \subset W$;
- ② compute $p' = P(\max_{w \in W_q} Z_w > t)$ with qMC quadrature;
- ③ estimate $\mathbb{P}_n(\max_{w \in W} Z_w > t)$ with

$$\hat{p} = p' + (1 - p') \hat{R}_q$$

where \hat{R}_q is a MC estimator of

$$R_q = \mathbb{P}_n \left(\max_{w \in W \setminus W_q} Z_w > t \mid \max_{w \in W_q} Z_w \leq t \right)$$

$\mathbb{P}_n(\max_{W \in W} Z_W > T)$: proposed hybrid algorithm

Algorithm:

- 1 select q grid points, denoted $W_q \subset W$;
- 2 compute $p' = P(\max_{w \in W_q} Z_w > t)$ with qMC quadrature;
- 3 estimate $\mathbb{P}_n(\max_{w \in W} Z_w > t)$ with


$$\hat{p} = p' + (1 - p')\hat{R}_q$$

where \hat{R}_q is a MC estimator of


$$R_q = \mathbb{P}_n \left(\max_{w \in W \setminus W_q} Z_w > t \mid \max_{w \in W_q} Z_w \leq t \right)$$

An asymmetric nested Monte Carlo scheme was developed for improved efficiency in R_q 's estimation. (See "orthant" paper and **anMC** R package).

Back to the test case with a conservative estimate...



Back to the test case with a conservative estimate...



NB: here, $\rho = 99.88829\%$ for a confidence of 99.12178% .

... and associated sequential strategies

For more on sequential conservative estimation



D. Azzimonti, D. Ginsbourger, C. Chevalier, J. Bect, Y. Richet (2021).

Adaptive Design of Experiments for Conservative Estimation of Excursion Sets.

Technometrics, 63:1, 13-26.

For more on sequential conservative estimation



D. Azzimonti, D. Ginsbourger, C. Chevalier, J. Bect, Y. Richet (2021).

Adaptive Design of Experiments for Conservative Estimation of Excursion Sets.

Technometrics, 63:1, 13-26.

Acknowledgements:

Drs Yann Richet and Grégory Caplin (French Nuclear Safety Institute) for providing the criticality safety test case.

Special thanks to Drs. Dario Azzimonti and Clément Chevalier for numerous invaluable inputs, and more generally, to all co-authors involved.

Outline

3 Miscellaneous

- Complements on Asymmetric Nested Monte Carlo
- Around profile extrema for excursion set visualization
- About Vorob'ev quantiles

Computation of the remainder

$$R_q = P_n(\max_{\mathbf{x} \in E \setminus E_q} Z(\mathbf{x}) > t \mid \max_{\mathbf{x} \in E_q} Z(\mathbf{x}) \leq t)$$

Standard Monte Carlo:


- ① draw realizations z_1^q, \dots, z_s^q from $Z(E_q) \mid \max_{\mathbf{x} \in E_q} Z(\mathbf{x}) \leq t$;
- ② for each z_i^q , draw a realization from $Z(E \setminus E_q) \mid Z(E_q) = z_i^q$;
- ③ Estimate R_q with $R_q^{\text{MC}} = \frac{1}{s} \sum_{i=1}^s \mathbf{1}_{\max(Z(E \setminus E_q)(\omega_i) \mid Z(E_q)=z_i^q) > t}$

The cost of step 1 is higher than the cost of step 2.

At fixed computational budget we reduce the variance of R_q^{MC} exploiting this difference with **asymmetric nested Monte Carlo**.

Computation of the remainder


At fixed computational budget we reduce the variance of R_q^{MC} drawing many realizations of $Z(E \setminus E_q) \mid Z(E_q) = z_i^q$ for each z_i .



Computation of the remainder

At fixed computational budget we reduce the variance of R_q^{MC} drawing many realizations of $Z(E \setminus E_q) \mid Z(E_q) = z_i^q$ for each z_i .

Asymmetric sampling scheme



Computation of the remainder: asymmetric nested MC

$$R_q = P_n(\max_{\mathbf{x} \in E \setminus E_q} Z(\mathbf{x}) > t \mid \max_{\mathbf{x} \in E_q} Z(\mathbf{x}) \leq t)$$

- ① draw realizations z_1^q, \dots, z_s^q from $Z(E_q) \mid \max_{\mathbf{x} \in E_q} Z(\mathbf{x}) \leq t$;
- ② for each z_i^q , draw $m^* > 1$ samples from $Z(E \setminus E_q) \mid Z(E_q) = z_i^q$;
- ③ $R_q^{\text{anMC}} = \frac{1}{s} \frac{1}{m^*} \sum_{i=1}^s \sum_{j=1}^{m^*} \mathbf{1}_{\max(Z(E \setminus E_q; \omega_{i,j}) \mid Z(E_q) = z_i^q) > t}$

$Var(R_q^{\text{anMC}})$ is optimally reduced if: $m^* = \sqrt{\frac{(\alpha+c)B}{\beta(A-B)}}$,

where $A = Var(\mathbf{1}_{\max(Z(E \setminus E_q) \mid Z(E_q)) > t})$,

$B = \mathbb{E}[Var(\mathbf{1}_{\max(Z(E \setminus E_q) \mid Z(E_q)) > t} \mid \max_{\mathbf{x} \in E_q} Z(\mathbf{x}) \leq t)]$ and α, β, c system dependent constants.

Azzimonti, D. and Ginsbourger D. (2018). *Estimating orthant probabilities of high dimensional Gaussian vectors with an application to set estimation*. *Journal of Computational and Graphical Statistics* 27:2, 255-267.

Properties of the anMC estimator

For each realization w_i we sample m realizations $y_{i,1}, \dots, y_{i,m}$ from $Y \mid w = W_i$, leading to a realization $\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m g(w_i, y_{i,j})$ of the estimator

$$\tilde{G} = \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m g(W_i, Y_{i,j}).$$

Proposition

Consider n independent copies W_1, \dots, W_n of W and, for each W_i , m copies $Y_{i,j} = Y_j \mid W_i$ $j = 1, \dots, m$, independent conditionally on W_i . Then,

$$\text{Var}(\tilde{G}) = \frac{1}{n} \text{Var}(g(W_1, Y_{1,1})) - \frac{m-1}{nm} \mathbb{E}[\text{Var}(g(W_1, Y_{1,1}) \mid W_1)].$$

Optimality for anMC

Under the same assumptions, \tilde{G} has minimal variance when

$$m = \tilde{m} = \sqrt{\frac{(\alpha + c)B}{\beta(A - B)}},$$


where $A = \text{Var}(g(W_1, Y_{1,1}))$ and $B = \mathbb{E}[\text{Var}(g(W_1, Y_{1,1}) | W_1)]$. Moreover denote with $\varepsilon = \tilde{m} - \lfloor \tilde{m} \rfloor$, then the optimal integer is $m^* = \lfloor \tilde{m} \rfloor$ if

$$\varepsilon < \frac{(2\tilde{m} + 1) - \sqrt{4(\tilde{m})^2 + 1}}{2}$$

or $m^* = \lceil \tilde{m} \rceil$ otherwise.



If $m^* > \frac{2(\alpha+c)B}{(c+\alpha)B+\beta(A-B)}$ then $\text{Var}(\tilde{G}) = \text{Var}(\hat{G})[1 - \eta]$, with $\eta \in (0, 1)$, where \hat{G} is the standard MC estimator.

Comparison GanMC/GMC with state-of-the-art




Benchmark: 6d GRF, discretization: 1000 Sobol' points, k Matérn ($\nu = 5/2$) with $\theta = [0.5, 0.5, 1, 1, 0.5, 0.5]^T$ and $\sigma^2 = 8$, m constant, $t = 5$.

More comparisons anMC/MC



Benchmark: 6d GRF, discretization: 1000 Sobol' points, k Matérn ($\nu = 5/2$) with $\theta = [0.5, 0.5, 1, 1, 0.5, 0.5]^T$ and $\sigma^2 = 8$, m constant.

Choice of m^* 

Benchmark: 6d GRF with m constant, k Matérn ($\nu = 5/2$), $\theta = [0.5, 0.5, 1, 1, 0.5, 0.5]$, $\sigma^2 = 8$. Discretized over first 1000 points of Sobol' sequence.

Quantity of interest:

$$1 - p = P(X < t), t = 5.$$

GanMC algorithm with p_q fixed, R_q (anMC) computed for different m .


Median estimated value $\hat{p} = 0.9644$,
 $\hat{p}_q = 0.9636$.

Outline

3 Miscellaneous

- Complements on Asymmetric Nested Monte Carlo
- Around profile extrema for excursion set visualization
- About Vorob'ev quantiles


Simulation of coastal flooding at “Les Boucholeurs”




Study site location (left) and computational domain limits (right, in white) with location of the forcing conditions (right, in blue).

Test case input and output parametrization

a: Forcing parameters (X)




b: Model results (inundation depth) and Y values



(a) Schematic representation of the tide and surge temporal signals and the different parameters describing them. (b) Maps of inland water height for given values of the parameters, and deduced value of flood surface.

Estimated coordinate profile maxima for the 5 inputs



Key underlying result

Theorem

Consider $(Z(\mathbf{x}))_{\mathbf{x} \in D} \sim GP(\mu, \mathfrak{K})$ and an approximating process of Z , \tilde{Z} , defined by $\tilde{Z}(\mathbf{x}) = \mathbf{a}(\mathbf{x}) + \mathbf{b}^T(\mathbf{x})Z_G$ where the \mathbf{a} , \mathbf{b} functions and $G = \{\mathbf{g}_1, \dots, \mathbf{g}_\ell\} \subset D$ ($\ell \geq 1$) are given. Then, for $T \subset D$ and any $u > \mu_T^{\tilde{\Delta}}$,

$$\mathbb{P} \left(\left| \sup_{\mathbf{x} \in T} Z(\mathbf{x}) - \sup_{\mathbf{x} \in T} \tilde{Z}(\mathbf{x}) \right| > u \right) \leq 2 \exp \left(- \frac{(u - \mu_T^{\tilde{\Delta}})^2}{2(\sigma_T^{\tilde{\Delta}})^2} \right), \quad (1)$$

where

$$\mu_T^{\tilde{\Delta}} = \sup_{\mathbf{x} \in T} |\mu^{\tilde{\Delta}}(\mathbf{x})| \text{ and } (\sigma_T^{\tilde{\Delta}})^2 = \sup_{\mathbf{x} \in T} \mathfrak{K}^{\tilde{\Delta}}(\mathbf{x}, \mathbf{x}) \text{ with} \quad (2)$$

$$\mu^{\tilde{\Delta}}(\mathbf{x}) = \mathbb{E}[Z(\mathbf{x}) - \tilde{Z}(\mathbf{x})] = \mu(\mathbf{x}) - \mathbf{a}(\mathbf{x}) - \mathbf{b}^T(\mathbf{x})\mu(G)$$

$$\mathfrak{K}^{\tilde{\Delta}}(\mathbf{x}, \mathbf{x}') = \mathfrak{K}(\mathbf{x}, \mathbf{x}') - \mathfrak{K}(\mathbf{x}', G)\mathbf{b}(\mathbf{x}) - \mathfrak{K}(\mathbf{x}, G)\mathbf{b}(\mathbf{x}') + \mathbf{b}^T(\mathbf{x})\mathfrak{K}(G, G)\mathbf{b}(\mathbf{x}'),$$

If $\tilde{Z} - Z$ is centred then (1) is valid for any $u > 0$.

For more detail

More on the [profile maxima approach and its application](#) to the BRGM data can be found in



D. Azzimonti, D. Ginsbourger, J. Rohmer, D. Idier (2019, to appear in Technometrics)

Profile extrema for visualizing and quantifying uncertainties on excursion regions. Application to coastal flooding.

<https://arxiv.org/abs/1710.00688>

For more detail

More on the [profile maxima approach and its application](#) to the BRGM data can be found in



D. Azzimonti, D. Ginsbourger, J. Rohmer, D. Idier (2019, to appear in Technometrics)

Profile extrema for visualizing and quantifying uncertainties on excursion regions. Application to coastal flooding.

<https://arxiv.org/abs/1710.00688>

For more on random fields and geometry, see in particular



R. J. Adler and J. E. Taylor (2007)

Random Fields and Geometry.

Springer

and references therein.

Outline

3 Miscellaneous

- Complements on Asymmetric Nested Monte Carlo
- Around profile extrema for excursion set visualization
- About Vorob'ev quantiles

Generalized optimality property for Vorob'ev quantiles

Proposition

For any $\rho \in [0, 1]$, the Vorob'ev quantile

$$Q_\rho = \{x \in D : p_n(x) \geq \rho\}$$

minimizes the expected distance in measure with Γ among measurable sets M such that $\mu(M) = \mu(Q_\rho)$, i.e.,

$$\mathbb{E}_n [\mu(Q_\rho \Delta \Gamma)] \leq \mathbb{E}_n [\mu(M \Delta \Gamma)],$$

for any measurable set M such that $\mu(M) = \mu(Q_\rho)$.

Generalized optimality property for Vorob'ev quantiles

Proposition

For any $\rho \in [0, 1]$, the Vorob'ev quantile

$$Q_\rho = \{x \in D : p_n(x) \geq \rho\}$$

minimizes the expected distance in measure with Γ among measurable sets M such that $\mu(M) = \mu(Q_\rho)$, i.e.,

$$\mathbb{E}_n [\mu(Q_\rho \Delta \Gamma)] \leq \mathbb{E}_n [\mu(M \Delta \Gamma)],$$

for any measurable set M such that $\mu(M) = \mu(Q_\rho)$.

A proof of this property is presented in Dario Azzimonti's PhD thesis (2016).